

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Siglec-1/CD169 in direct ELISAs.
<b>Source</b>	Recombinant Monoclonal Mouse IgG <sub>1</sub> Clone # 908102
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human Siglec-1/CD169 Ser20-Gln1641 Accession # Q9BZZ2
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	0.25 µg/mL	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

Siglecs are sialic acid specific I-type lectins that belong to the immunoglobulin superfamily. Structurally, they are transmembrane proteins with an N-terminal Ig-like V-set domain followed by varying numbers of Ig-like C2-set domains (1, 2). Human Siglec-1, also known as sialoadhesin and CD169, is a 175-185 kDa glycoprotein. It contains a 1622 amino acid (aa) extracellular domain (ECD) with one Ig-like V-set domain and 16 Ig-like C2-set domains, a 21 aa transmembrane segment, and a 44 aa cytoplasmic domain (3). Within the ECD, human Siglec-1 shares approximately 70% aa sequence identity with mouse and rat Siglec-1. Alternate splicing generates a potentially soluble form of the ECD, and a second isoform with a substituted cytoplasmic domain. Siglec-1 expression is restricted to lymph node and splenic macrophages, plus some tissue macrophages (3). The adhesive function of Siglec-1 is supported by the N-terminal Ig-like domain which shows a selectivity for  $\alpha$ 2,3-linked sialic acid residues (3-5). Siglec-1 binds a number of sialylated molecules including the mannose receptor, MGL1, MUC1, PSGL-1, and different glycoforms of CD43 (6-9). Its binding capacity can be masked by endogenous sialylated molecules (10, 11). The sialylated and sulfated N-linked carbohydrates that modify Siglec-1 itself are required for ligand binding (6, 7). Siglec-1 is expressed on dendritic cells following rhinovirus exposure, and these DC promote T cell energy (12). It is also induced on circulating monocytes during systemic sclerosis and HIV-1 infection (13-15). Siglec-1 can trap HIV-1 particles for *trans* infection of permissive cells (14).

**References:**

1. Varki, A. and T. Angata (2006) *Glycobiology* **16**:1R.
2. Crocker, P.R. *et al.* (2007) *Nat. Rev. Immunol.* **7**:255.
3. Hartnell, A. *et al.* (2001) *Blood* **97**:288.
4. Nath, D. *et al.* (1995) *J. Biol. Chem.* **270**:26184.
5. Crocker, P.R. *et al.* (1991) *EMBO J.* **10**:1661.
6. Martinez-Pomares, L. *et al.* (1999) *J. Biol. Chem.* **274**:35211.
7. Kumamoto, Y. *et al.* (2004) *J. Biol. Chem.* **279**:49274.
8. Nath, D. *et al.* (1999) *Immunology* **98**:213.
9. van den Berg, T.K. *et al.* (2001) *J. Immunol.* **166**:3637.
10. Nakamura, K. *et al.* (2002) *Glycobiology* **12**:209.
11. Barnes, Y.C. *et al.* (1999) *Blood* **93**:1245.
12. Kirchberger, S. *et al.* (2005) *J. Immunol.* **175**:1145.
13. York, M.R. *et al.* (2007) *Arthritis Rheum.* **56**:1010.
14. Rempel, H. *et al.* (2008) *PLoS ONE* **3**:e1967.
15. van der Kuyl, A.C. *et al.* (2007) *PLoS ONE* **2**:e257.