

DESCRIPTION


| | |
|---------------------------|---|
| Species Reactivity | Human |
| Specificity | Detects recombinant human IL-34 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant mouse IL-34 is observed. |
| Source | Monoclonal Mouse IgG ₁ Clone # 578416 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | Mouse myeloma cell line NS0-derived recombinant human IL-34 Asn21-Pro242 Accession # Q6ZMJ4 |
| Endotoxin Level | <1.0 EU per 1 µg of the antibody by the LAL method. |
| Formulation | Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|---|---|--|
| Intracellular Staining by Flow Cytometry | 2.5 µg/10 ⁶ cells | Human IL-34 transfected NS0 cells fixed with paraformaldehyde and permeabilized with saponin |
| CyTOF-ready | Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation. | |
| Neutralization | Measured by its ability to neutralize IL-34-induced proliferation in human peripheral blood monocytes. Lin, H. <i>et al.</i> (2008) <i>Science</i> 320 :807. The Neutralization Dose (ND ₅₀) is typically 0.01-0.05 µg/mL in the presence of 10 ng/mL Recombinant Human IL-34. | |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Reconstitution | Reconstitute at 0.5 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

Interleukin 34 (IL-34; also known as uncharacterized protein C16orf77) is secreted as a homodimer consisting of 39 kDa monomers (1). It belongs to no known cytokine family. Human IL-34 is synthesized as a 242 amino acid (aa) precursor that contains a 20 aa signal sequence and a 222 aa mature chain. The mature chain contains one potential site of N-linked glycosylation. Human IL-34 is 71% identical to mouse IL-34 on the amino acid level (1). IL-34 is expressed in various tissues, including the heart, brain, liver, kidney, spleen, thymus, testes, ovary, small intestine, prostate, and colon, and is most abundant in the spleen (1). The receptor for IL-34 is colony-stimulating factor 1 receptor (CSF-1R) (1). IL-34 stimulates monocyte proliferation (1). In functional studies, IL-34, like CSF-1, the other ligand for CSF-1R, stimulated phosphorylation of extracellular signal-regulated kinase-1 and -2 (ERK1/2) in human monocytes (1). In addition, IL-34 promoted the formation of the colony-forming unit-macrophage (CFU-M), a macrophage progenitor, in human bone marrow cultures (1).

References:

1. Lin, H. *et al.* (2008) *Science* **320**:807.