

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse Notch-1 intracellular domain in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # N1A
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse embryonic fibroblast cell line PA317-derived recombinant mouse Notch-1 Accession # Q01705
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	0.25 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Notch-1 is a 300 kDa type I transmembrane glycoprotein that is one of four Notch homologues involved in developmental processes (1-3). Notch signaling is important for maintaining stem cells and inducing differentiation, especially in the nervous system and lymphoid tissues (2-4). Notch can specify binary cell fates. For example, it promotes T cell over B cell development from a common precursor (2). Mouse Notch-1 is synthesized as a 2531 amino acid (aa) precursor that contains an 18 aa signal sequence, a 1707 aa extracellular domain (ECD) with 36 EGF-like repeats and three Lin-12/notch repeats (LNR), a 21 aa transmembrane (TM) segment and a 785 aa cytoplasmic domain that contains six ankyrin repeats, a glutamine-rich domain and a PEST sequence. The 11th and 12th EGF-like repeats, that bind ligands such as Jagged and Delta-like families in humans, correspond to aa 412-488 in mouse Notch-1 (6). Elongation of O-linked fucose chains by Fringe family members at a site within this region can inhibit the interaction of Notch with Jagged ligands, thereby promoting Delta-like ligand interactions (7). The Notch-1 receptor undergoes post-translational furin-type proteolytic cleavage, generating a heterodimer through the interaction of a hydrophobic area C-terminal to the LNR on the extracellular region with the transmembrane/cytoplasmic portion (8, 9). Upon ligand binding, additional sequential proteolysis by TNF-converting enzyme (ADAM17) and the presenilin-dependent γ -secretase results in the release of the Notch intracellular domain (NICD) which translocates into the nucleus, activating transcription of Notch-responsive genes (10).

References:

1. Ellisen, L.W. *et al.* (1991) *Cell* **66**:649.
2. Dumortier, A. *et al.* (2005) *Int. J. Hematol.* **82**:277.
3. Yoon, K. and N. Gaiano (2005) *Nat. Neurosci.* **8**:709.
4. Androutsellis-Theotokis, A. *et al.* (2006) *Nature* **442**:823.
5. Weng, A.P. *et al.* (2004) *Science* **306**:269.
6. Hambleton, S. *et al.* (2004) *Structure* **12**:2173.
7. Yang, L. *et al.* (2005) *Mol. Biol. Cell* **16**:927.
8. Sanchez-Irizarry, C. *et al.* (2004) *Mol. Cell. Biol.* **24**:9265.
9. Logeat, F. *et al.* (1998) *Proc. Natl. Acad. Sci. USA* **95**:8108.
10. Mumm, J.S. and R. Kopan (2000) *Dev. Biol.* **228**:151.