

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human DAPK3/ZIPK in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) DAPK1, rhDAPK2, rhDRAK1, or rhDRAK2 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 729830
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human DAPK3/ZIPK Arg347-Arg454 Accession # O43293
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.5 µg/mL	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.5 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

DAPK3 (Death-associated protein kinase 3; also ZIP kinase and DLK) is a 52 kDa member of the CAMK Ser/Thr kinase family, protein kinase superfamily of enzymes. In general, mouse DAPK3 is nuclear while human DAPK3 is cytosolic. DAPK3 is found in smooth muscle where it phosphorylates the regulatory light chain of myosin and stabilizes thick filament formation. Human DAPK3 is 454 amino acids (aa) in length. It contains one protein kinase domain (aa 134-275) and a C-terminal Leu-zipper motif (aa 427-441) that mediates homo- and hetero-dimerization. There is one 35 kDa alternate splice form that shows a 10 aa substitution for aa 313-454. This form has both a nuclear and cytosolic localization. Over aa 347-454, human DAPK3 is 65% aa identical to mouse DAPK3.