

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Nicastrin in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 716918
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Nicastrin Asn34-Glu669 Accession # Q92542
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Immunocytochemistry	8-25 µg/mL	See Below

DATA

Immunocytochemistry



Nicastrin in HepG2 Human Cell Line.
Nicastrin was detected in immersion fixed HepG2 human hepatocellular carcinoma cell line using Mouse Anti-Human Nicastrin Monoclonal Antibody (Catalog # MAB5378) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). Specific staining was localized to cytoplasm. View our protocol for [Fluorescent ICC Staining of Cells on Coverslips](#).

PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Nicastrin (NCT) is a 150-160 kDa member of the Nicastrin family of proteins. It is a component of the aspartyl protease γ-secretase complex and serves to stabilize and direct γ-secretase components to proper positions in the plasma membrane. The γ-secretase complex mediates the cleavage of intramembrane proteins such as Notch-1 and APP. Mature human Nicastrin is a 676 amino acid type I transmembrane glycoprotein. It contains a 636 aa extracellular domain (aa 34-669) that shows a 58 aa sequence (aa 312-369) which interacts with γ-secretase substrates. There are multiple splice variants of NCT. One shows a deletion of aa 195-322 and 394-709, a second shows a 29 aa substitution for the C-terminal 604 aa and a third shows a deletion of aa 200-709 accompanied by an insertion of 33 aa after Leu30. Over aa 34-669, human NCT shares 90% aa identity with mouse NCT.