

## DESCRIPTION

<b>Species Reactivity</b>	Rat
<b>Specificity</b>	Detects rat EphB1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human EphA1, recombinant mouse EphA2, A3, A4, A6, A7, A8, B2, B3, B4, B6, or recombinant rat EphA5 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 88506
<b>Purification</b>	Protein A or G purified from ascites
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant rat EphB1 Met18-Gln538 Accession # P09759
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	Recombinant Rat EphB1 Fc Chimera (Catalog # <a href="#">1596-B1</a> )

## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

EphB1, also known as Elk, Cek6, Net, and Hek6 (1), is a member of the Eph receptor family which binds members of the ephrin ligand family. There are two classes of receptors, designated A and B. Both the A and B class receptors have an extracellular region consisting of a globular domain, a cysteine-rich domain, and two fibronectin type III domains. This is followed by the transmembrane region and the cytoplasmic region. The cytoplasmic region contains a juxtamembrane motif with two tyrosine residues, which are the major autophosphorylation sites, a kinase domain, and a conserved sterile alpha motif (SAM) in the carboxy tail which contains one conserved tyrosine residue. Activation of kinase activity occurs after ligand recognition and binding. EphB1 has been shown to bind ephrin-B2, ephrin-B1, ephrin-A3, ephrin-A1, ephrin-A4, and ephrin-B3 (2, 3). The extracellular domains of human and rat EphB1 share 99% amino acid identity. Only membrane-bound or Fc-clustered ligands are capable of activating the receptor *in vitro*. While soluble monomeric ligands bind the receptor, they do not induce receptor autophosphorylation and activation (2). *In vivo*, the ligands and receptors display reciprocal expression (3). It has been found that nearly all the receptors and ligands are expressed in developing and adult neural tissue (3). The ephrin/Eph families also appear to play a role in angiogenesis (3).

## References:

1. *Eph Nomenclature Committee [letter]*. (1997) Cell **90**:403.
2. Flanagan, J.G. and P. Vanderhaeghen (1998) Annu. Rev. Neurosci. **21**:309.
3. Pasquale, E.B. (1997) Curr. Opin. Cell Biol. **9**:608.