

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Complement Factor H-related 2/CFHR2 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) CFHR1 or rhCFHR5 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 510511
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Complement Factor H-related 2/CFHR2 Glu19-Glu268 Accession # P36980
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	Recombinant Human Complement Factor H-related 2/CFHR2
Immunoprecipitation	25 µg/mL	Conditioned cell culture medium spiked with Recombinant Human Complement Factor H-related 2/CFHR2, see our available Western blot detection antibodies

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Complement Factor H-related 2 (CFHR2) is synthesized in the liver and secreted into plasma. It may be involved in complement regulation. It can also be associated with lipoproteins and may play a role in lipid metabolism. Human CFHR2 shares 48% identity with that of rat.