

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse IL-15 R α in direct ELISAs.
Source	Monoclonal Rat IgG _{2B} Clone # 888220
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse IL-15 R α Gly33-Lys205 Accession # Q60819
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25 μ g/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Interleukin 15 Receptor alpha (IL-15 R α) is a high affinity receptor that specifically binds IL-15 with high affinity and associates as a heterotrimer with the IL-2 receptors beta and gamma subunits to initiate signal transduction. IL-15 R α is expressed on a wide variety of T cells and B cells as well as non-lymphoid cells. IL-15 R α is a 58-60 kDa protein that shares structural similarities to the IL-2 R α protein. IL-15 R α and IL-2 R α genes also share similar intron-exon organization and are closely linked on human chromosome 10p14-p15. Human IL-15 R α shares 45% amino acid (aa) homology with the mouse form of the receptor. Signaling of IL-15 can occur in one of three ways; through the heterotrimeric complex of IL-15 R α , IL-2 R β and IL-2 R γ c, through the heterodimeric complex of IL-2 receptors beta and gamma common, through a novel 60-65 kDa IL-15 RX subunit found on mast cells. The binding of IL-15 to IL-15 R α has been reported to antagonize the TNF- α -mediated apoptosis in fibroblasts by competing with TNFRI for TRAF2 binding.