RDsystems a biotechne brand

Human CA125/MUC16 Antibody

Monoclonal Mouse IgG2A Clone # 986808 Catalog Number: MAB5609

DESCRIPTION		
Species Reactivity	Human	
Specificity	Detects human CA125/MUC-16 in direct ELISAs.	
Source	Monoclonal Mouse IgG _{2A} Clone # 986808	
Purification	Protein A or G purified from ascites	
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human CA125/MUC-16 Met13360-Gln14347 (Met13472Thr & Gln13957Lys) Accession # NP_078966.2	
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended	Sample
	Concentration	
Flow Cytometry	0.25 μL/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	



- *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C Stability & Storage
 - Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
 - 12 months from date of receipt, -20 to -70 °C as supplied.
 - 1 month, 2 to 8 °C under sterile conditions after reconstitution. ٠
 - 6 months, -20 to -70 °C under sterile conditions after reconstitution.

Rev. 7/21/2022 Page 1 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449



Human CA125/MUC16 Antibody

Monoclonal Mouse IgG_{2A} Clone # 986808 Catalog Number: MAB5609

BACKGROUND

MUC16, also known as the CA125 antigen, is a mucin protein that may be found in type I transmembrane or secreted forms that are used monitor the progress of epithelial ovarian cancer therapy (1, 2). Expression of isoforms, proteolytic cleavage, and heavy N- and O- linked glycosylation produce forms of human MUC16 that can vary from 1148 to 22152 amino acids (aa) in length and 200 - 5000 kDa in size (1, 2). The 22152 aa form contains ser/thr-rich N-terminal tandem repeats, 4 LRR (Leu-rich repeat) domains, 56 SEA (sea urchin sperm protein, enterokinase and agrin) domains, a transmembrane domain, and a 31 aa cytoplasmic domain that includes a tyrosine phosphorylation site (1-4). SEA domains are ~120 aa in length, contain conserved residues including potential O-glycosylation sites and a pair of cysteines, and are often found in transmembrane mucins (3). The protein produced by R&D Systems represents aa 13360-14347 of the full sequence and includes the last 6 SEA domains. It shares 68% aa identity with canine MUC16. MUC16 is over-expressed by tumor cells including ovarian and mesothelial cancers (5). The transmembrane form can adhere to mesothelin in the peritoneum, facilitating metastasis of ovarian cancer to the peritoneal cavity (5-7). MUC16 also binds galectin-1 on immune cells and enhances its expression on tumor cells (8). MUC16-expressing tumors adhere to NK cells, down-regulate CD16 and suppress NK response, which may promote immune evasion (9, 10). MUC16 is also cyclically expressed in the endometrium and may contribute to immune privilege during pregnancy (10). In the eye, MUC16 and other mucins protect the cornea and keep it hydrated. It is altered on the conjunctival epithelium of patients with non-Sjogren dry eye syndrome (11).

References:

- 1. Yin, B. W. T. K. O. Lloyd, 2001, J. Biol. Chem. 276:27371.
- 2. Maeda, T. et al. (2004) J. Biol. Chem. 279:13174.
- 3. Fendrick, J. L. et al. (1997) Tumour Biol. 18:278.
- 4. Swissprot accession Q8WXI7.
- 5. Kaneko, O. et al. (2009) J. Biol. Chem. 284:3739.
- 6. Rump, A. et al. (2004) J. Biol. Chem. 279:9190.
- 7. Gubbels, J. A. A. et al. (2006) Mol. Cancer 5:50.
- 8. Seelenmeyer, C. et al. (2003) J. Cell Sci. 116:1305.
- 9. Patankar, M. S. et al. (2005) Gynecol. Oncol. 99:704.
- 10. Belisle, J. A. et al. (2007) Immunology 122:418.
- 11. Blalock, T. D. et al. (2007) Invest. Ophthalmol. Vis. Sci. 48:4509.

Rev. 7/21/2022 Page 2 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449