

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human TRF2 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human TRF1 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 689524
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human TRF2 Gln78-Thr238 (predicted) Accession # Q15554
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.5 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

TRF2 (telomeric repeat-binding factor 2; also telomeric DNA-binding protein) is a 55-60 kDa, ubiquitously expressed nuclear protein that participates in telomere homeostasis. TRF-2 binds as a dimer to TTAGGG repeats at ends of chromosomes (telomeres), where it blocks inappropriate activation of the ATM/p53 pathway. It also collaborates with TRF1 to promote normal telomere length. Human TRF2 is 500 amino acids (aa) in length. It contains an N-terminal Arg-rich region (aa 13-30), a dimerization domain (aa 46-112), an NLS (aa 329-333), and a DNA binding HTH myb-type domain (aa 442-499). There is one potential alternate start site 42 aa upstream of the standard start site, and one splice form that shows a 13 aa substitution for aa 239-500. Over aa 78-238, human TRF2 shares 96% aa identity with mouse TRF2.