

**DESCRIPTION**

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human IL-411 in direct ELISAs.
<b>Source</b>	Monoclonal Rat IgG <sub>2B</sub> Clone # 1006209
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human IL-411 Met1-His567 Accession # Q96RQ9
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

<b>ELISA</b>	This antibody functions as an ELISA detection antibody when paired with Rat Anti-Human IL-411 Monoclonal Antibody (Catalog # MAB56842).  <i>This product is intended for assay development on various assay platforms requiring antibody pairs.</i>
--------------	---

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

Interleukin 4 induced protein 1 (IL-411), also known as protein FIG-1 and L-amino acid oxidase, is encoded by a B-cell IL-4-inducible gene, FIG1, and is highly expressed in primary metastinal B-cell lymphomas (1-4). It belongs to the flavin monoamine oxidase family, FIG1 subfamily. Enzymological characterization reveals that IL-411 has L-amino acid oxidase activity with preference toward aromatic amino acids. Studies have shown that hIL-411 inhibited the proliferation of CD3-stimulated T lymphocytes with a similar effect on CD4(+) and CD8(+) T cells (5). Its inhibitory effect was dependent on enzymatic activity and H<sub>2</sub>O<sub>2</sub> production. Its restricted expression to lymphoid tissues indicates that it may play an important function in the immune system (1, 4).

**References:**

1. Chu, C.C. and W.E. Paul. (1997) Proc. Natl. Acad. Sci. USA **94**:2507.
2. Mason, J.M. *et al.* (2004) J. Immunol. **173**:4561.
3. Chavan, S.S. *et al.* (2002) Biochim. Biophys. Acta. **1576**:70.
4. Copie-Bergman, C. *et al.* (2003) Blood **101**:2756.
5. Boulland, M.L. *et al.* (2007) Blood **110**:220.