

DESCRIPTION


Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse and rat Thioredoxin-2 in Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 543317
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Thioredoxin-2 Thr60-Gly166 Accession # Q99757
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 µg/mL	See Below
Simple Western	5 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Thioredoxin-2 (Trx-2; also known as mitochondrial Thioredoxin) belongs to the evolutionarily conserved Thioredoxin family of proteins. Thioredoxins share the Thioredoxin fold containing the active site-CGPC motif. In their reduced form, the active site cysteine residues reduce protein disulfides. The resulting active site disulfide is subsequently reduced in a reaction catalyzed by a NADPH-dependent Thioredoxin reductase. Thioredoxin-2 contains an N-terminal 59 amino acid (aa) transit sequence that is cleaved upon translocation to mitochondria. The amino acid sequence of mature human Thioredoxin-2 is 98% identical to mouse and rat Thioredoxin-2. Thioredoxin-2 interacts with specific components of the mitochondrial respiratory chain and helps regulate the membrane potential. Thioredoxin-2 is ubiquitously but variably expressed and high expression confers resistance to oxidant-induced apoptosis.