

DESCRIPTION


Species Reactivity	Human/Equine
Specificity	Detects recombinant human 5'-Nucleotidase in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant mouse 5'-Nucleotidase is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 606112
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human 5'-Nucleotidase/CD73 Trp27-Ile511 Accession # P21589
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CytoF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

5'-Nucleotidase (also [ecto]-5'-nucleotidase/5'-NT, designated CD73) is a variably glycosylated, 69-73 kDa member of the 5'-Nucleotidase family of enzymes. It is expressed on multiple cell types, including vascular endothelium, transitional and nonkeratinized epithelium, cardiomyocytes, small intestine epithelium, FoxP3⁺ Treg lymphocytes, FDCs and B cells. 5'-Nucleotidase hydrolyzes AMP to adenosine and phosphate. This creates diffusible nucleosides necessary for cell homeostasis, and a ligand for cell membrane adenosine receptors. Mature human 5'-Nucleotidase is a 523 amino acid (aa) GPI-linked protein (aa 27-549). It contains a large Zn-dependent nucleotidase catalytic region (aa 28-532) and a C-terminal substrate binding site (aa 500-506). On the cell surface it exists as a disulfide-linked homodimer. Two splice variants are reported. One shows a deletion of aa 405-454, and a second possesses a 12 aa substitution for aa 253-574. Over aa 1-511, human 5'-Nucleotidase shares 88% aa identity with both mouse and rat 5'-Nucleotidase.