

## DESCRIPTION

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse BMP-15/GDF-9B in direct ELISAs. In this format, no cross-reactivity with recombinant human BMP1.1, 2, 3, 3b, 4, 5, 6, 7, 8, 9, 10, 15, recombinant mouse BMP-3, 3b, 4, 5, 6, 7, 8b, 9, or 10 is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 582703
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant mouse BMP-15/GDF-9B Gln268-Arg392 Accession # Q9Z0L4
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.


## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Immunohistochemistry</b>	8-25 µg/mL	See Below

## DATA

### Immunohistochemistry



**BMP-15/GDF-9B in Mouse Ovary.** BMP-15/GDF-9B was detected in perfusion fixed frozen sections of adult mouse ovary using Mouse BMP-15/GDF-9B Monoclonal Antibody (Catalog # MAB5917) at 10 µg/mL overnight at 4 °C. Tissue was stained using the NorthernLights™ 557-conjugated Anti-Rat IgG Secondary Antibody (red; Catalog # Catalog # NL013) and counterstained with DAPI (blue). Specific staining was localized to an oocyte within the ovary. View our protocol for [Fluorescent IHC Staining of Frozen Tissue Sections](#).

## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Bone morphogenetic protein 15 (BMP-15), also known as GDF-9B, is a member of the TGF-β superfamily. Mature BMP-15 has three intrachain disulfide bands that form a cysteine-knot fold. It is secreted as a 34 kDa non-disulfide-linked homodimer and as a 37 kDa nondisulfide heterodimer with GDF-9. BMP-15 is a product of oocytes and promotes granulosa cell proliferation and stem cell factor secretion. Mature mouse BMP-15 shares 44% aa sequence identity with mature mouse GDF-9. It also shares 70% and 78% aa sequence identity with human and sheep BMP-15, respectively.