

DESCRIPTION


| | |
|---------------------------|---|
| Species Reactivity | Human |
| Specificity | Detects human UCH-L3 in direct ELISAs. In direct ELISAs and Western blots, no cross-reactivity with recombinant human UCH-L1 is observed. |
| Source | Monoclonal Mouse IgG ₁ Clone # 677710 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | <i>E. coli</i> -derived recombinant human UCH-L3 Glu2-Ala230 Accession # P15374 |
| Formulation | Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|----------------------------|----------------------------------|--|
| Western Blot | 3 µg/mL | See Below |
| Immunoprecipitation | 25 µg/mL | Cell lysates spiked with Recombinant Human UCH-L3, see our available Western blot detection antibodies |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Reconstitution | Sterile PBS to a final concentration of 0.5 mg/mL. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

UCH-L3 (Ubiquitin carboxyterminal hydrolase isozyme 3) is a 26-28 kDa member of the peptidase C12 family of enzymes. It shows wide expression, being noted in adipocytes, renal duct epithelium, neurons, and striated muscle. UCH-L3 cleaves both monomeric ubiquitin (Ub) from Ub-protein conjugates, and a GGLRQ peptide from the C-terminus of the Ub-like protein NEDD-8. Notably, UCH-L3 activity is muted in the presence of Ub interacting dimers. Human UCH-L3 is 230 amino acids (aa) in length. It is phosphorylated on Ser75 and Ser130 and contains two ubiquitin-binding sequences between aa 40-57 and 178-186. Four potential splice forms have been reported. Two show a four aa deletion between aa 15-18, with one also containing a 14 aa substitution for aa 179-230. Two others contain an alternative start site at Met37, one of which is also accompanied by a seven aa substitution for aa 143-230. Full length human UCH-L3 shares 98% aa identity with mouse and rat UCH-L3.