

**DESCRIPTION**


<b>Species Reactivity</b>	Human/Mouse
<b>Specificity</b>	Detects human and mouse PKN1 in Western blots.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 540421
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human PKN1 Ile215-Thr388 Accession # Q16512
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

PKN1 (serine/threonine protein kinase N1), also known as PKN or PRK, is a ubiquitous 120 kDa cytoplasmic kinase of the PKC superfamily. The 942 amino acid (aa) PKN1 contains three HR1 repeats (aa 34-263), a C2 domain (aa 325-461), and a protein kinase domain (aa 615-874). PKN1 can be activated by proteolysis, lipid interaction, or Rho G proteins, and is a downstream effector of RhoA. It promotes the activation of several kinases involved in regulation of the actin cytoskeleton, cell cycle, and apoptosis. In vivo, PKN1 phosphorylates histone H3 at the Thr11 position.