

**DESCRIPTION**


<b>Species Reactivity</b>	Human/Mouse
<b>Specificity</b>	Detects human and mouse ROR $\gamma$ /RORC/NR1F3 in Western blots.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 600214
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human ROR $\gamma$ /RORC/NR1F3 Met1-Gln100 Accession # P51449-2
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 $\mu$ m filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.5 $\mu$ g/mL	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

ROR $\gamma$  (Nuclear receptor ROR-gamma), also called RORC and NR1F3, is a member of the NR1 nuclear hormone receptor family. ROR $\gamma$  is a DNA binding transcription factor. ROR $\gamma$  is 518 amino acids (aa) in length. Deletion in mice implicates ROR $\gamma$  as being essential for lymphoid organogenesis and controlling apoptosis during thymopoiesis. Two splice forms differing in the first 24 aa have been found for this gene; isoform 2 deletes aa 1-21 resulting in an alteration of aa 22-24. Over aa 1-100 human RORC shares 96% identity with mouse ROR $\gamma$ .