

## DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects recombinant human ROR $\gamma$ /RORC2/NR1F3 by direct ELISAs.
<b>Source</b>	Recombinant Monoclonal Rabbit IgG Clone # 1181A
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	<i>E. coli</i> derived recombinant human ROR $\gamma$ /RORC2/NR1F3 Met1-Gln100 Accession # P51449
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 $\mu$ m filtered solution in PBS.


## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Intracellular Staining by Flow Cytometry</b>	0.25 $\mu$ g/10 <sup>6</sup> cells	See Below


## DATA

### Intracellular Staining by Flow Cytometry



**Detection of ROR $\gamma$ /RORC2/NR1F3 in Human PBMCs by Flow Cytometry.**  
Human peripheral blood mononuclear cells (PBMCs) either (A) unstimulated or (B) stimulated to induce Th17 cells were stained with Rabbit Anti-Human ROR $\gamma$ /RORC2/NR1F3 Monoclonal Antibody (Catalog # MAB61093) followed by Phycoerythrin-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog # F0110) and Mouse Anti-Human CD4 APC-conjugated Monoclonal Antibody (Catalog # FAB3791A). Quadrant markers were set based on control antibody staining (Catalog # MAB1050). To facilitate intracellular staining, cells were fixed and permeabilized with FlowX FoxP3 Fixation & Permeabilization Buffer Kit (Catalog # FC012). View our protocol for [Staining Intracellular Molecules](#).

### Intracellular Staining by Flow Cytometry



**Detection of ROR $\gamma$ /RORC2/NR1F3 in HepG2 cells by Flow Cytometry.**  
HepG2 cells were stained with Rabbit Anti-Human ROR $\gamma$ /RORC2/NR1F3 Monoclonal Antibody (Catalog # MAB61093, filled histogram) or isotype control antibody (Catalog # AB-105-C, open histogram), followed by Allophycocyanin-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog # F0111). To facilitate intracellular staining, cells were fixed and permeabilized with FlowX FoxP3 Fixation & Permeabilization Buffer Kit (Catalog # FC012). View our protocol for [Staining Intracellular Molecules](#).

## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Retinoic acid-related Orphan Receptor gamma (ROR $\gamma$ , TOR, RORC; NR1F3) is a member of the orphan nuclear receptor family. ROR $\gamma$  is expressed in the muscle, thymus, testis, pancreas, prostate, heart, and liver. ROR $\gamma$  plays a role in thymocyte development and homeostasis. RORs bind to DNA as monomers on half-site elements with 5' A/T-rich extensions. An N-terminal isoform of ROR $\gamma$ , ROR $\gamma$ t, has been shown to be specifically expressed in the thymus.