

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human LAT in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 661002
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human LAT His30-Gln113 Accession # O43561
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL. For liquid material, refer to CoA for concentration.
Shipping	Lyophilized product is shipped at ambient temperature. Liquid small pack size (-SP) is shipped with polar packs. Upon receipt, store immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

LAT (Linker for activation of T cells; also pp36) is a 36-38 kDa type III (no signal sequence; extracellular N-terminus) transmembrane adaptor phosphoprotein. It is expressed on mast cells, NK cells, monocytes (but not macrophages), platelets and T cells. LAT is phosphorylated upon ligation of the TCR, GPVI, and Fc epsilon RI, and serves as an anchor for several SH2-domain containing signaling proteins, including PLCγ1, SLP76 and Grb2. Human LAT is 262 amino acids (aa) in length. It contains a four aa extracellular segment with a 235 aa cytoplasmic domain (aa 28-262). Palmitoylation occurs on Cys26 and 29. There are 13 Ser/Thr phosphorylation sites plus five Tyr sites. Multiple potential splice variants are reported. There is an alternative start site 36 aa upstream of the standard site that may be accompanied by a deletion of aa 114-142. There is also a deletion of aa 88-97, and three substitutions: two aa for aa 215-262, 38 aa for aa 55-262, and 50 aa for aa 35-262. Over aa 30-113, human LAT shares 73% aa identity with mouse LAT.