

## DESCRIPTION


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	This antibody detects mouse Complement Component C5a R1 in direct ELISAs.
<b>Source</b>	Monoclonal Rat IgG <sub>2B</sub> Clone # 583837
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	NS0 mouse myeloma cell line transfected with mouse Complement Component C5a R1 Accession # P30993
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.5 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

C5a R1, also known as C5a ligand and CD88, is a 7TM protein expressed on myeloid, endothelial, epithelial, and smooth muscle cells. C5a R1 binds the activated complement anaphylatoxin C5a. In established allergic environments, this triggers neutrophil and eosinophil chemotaxis and the release of proinflammatory mediators. In contrast, C5a R1/C5a interactions are protective during allergen sensitization. Mouse C5a R1 shares 66% and 77% amino acid sequence identity with human and rat C5a R, respectively.