

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human Aggrecan in direct ELISAs. It specifically recognizes the N-terminal fragment of recombinant human Aggrecan released following cleavage by ADAMTS5.
Source	Monoclonal Mouse IgG _{2B} Clone # 725913
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	KLHG-conjugated human Aggrecan synthetic peptide Leu384-Glu392 Accession # P16112
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	2 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Aggrecan, together with type II collagen, makes up to 90% of the dry weight of healthy cartilage. It hydrates the collagen network and thus provides cartilage with its properties of compressibility and elasticity. The N-terminal region of human Aggrecan (residues 20-675) consists of two globular domains (G1 and G2), flanking an interglobular domain (IGD). The IGD contains the cleavage sites for matrix metalloproteinases and aggrecanases.