

Human TSH R Antibody

Monoclonal Mouse IgG₁ Clone # 484404 Catalog Number: MAB65342


DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human TSH R in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 484404
Purification	Protein A or G purified from ascites
Immunogen	NS0 mouse myeloma cell line transfected with human TSH R Accession # P16473
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25 μg/10 ⁶ cells	See Below

DATA



Detection of TSH R in HEK293 Human Cell Line Transfected with Human TSH R and eGFP by Flow Cytometry. HEK293 human embryonic kidney cell line transfected with either (A) human TSH R or (B) irrelevant transfectants and eGFP was stained with Mouse Anti-Human TSH R Monoclonal Antibody (Catalog # MAB65342) followed by Allophycocyanin-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # F0101B), Quadrant markers were set based on control antibody staining (Catalog # MAB002). View our protocol for Staining Membrane-associated Proteins.

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
	*Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
	 12 months from date of receipt, -20 to -70 °C as supplied.
	 1 month, 2 to 8 °C under sterile conditions after reconstitution.
	 6 months20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Thyroid stimulating hormone receptor (TSH R; also thyrotropin receptor) is an 85 kDa (unglycosylated) member of the G-protein coupled receptor 1 family. Human TSH R is synthesized as a 764 amino acid (aa) precursor that contains a 20 aa signal sequence, a 393 aa extracellular domain (ECD), a 269 aa membrane spanning domain consisting of seven transmembrane segments, and an 82 aa cytoplasmic domain. The ECD contains six leucine-rich repeats (LRR) and six potential sites of N-linked glycosylation. There are two isoforms, long and short, produced by a splicing variant corresponding to aa 253-764 in the long isoform that is missing in the short isoform. Human TSH R is 86-87% aa identical to mouse and rat TSH R. TSH R is the receptor for thyroid stimulating hormone, which plays a central role in controlling thyroid cell metabolism.

Rev. 2/7/2018 Page 1 of 1

