

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human FSH R in direct ELISAs.
Source	Monoclonal Mouse IgG _{2A} Clone # 626717
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with human FSH R Accession # M65085
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Neutralization	Measured by its ability to neutralize FSH α/β-induced cAMP production in the HEK293 human embryonic kidney cell line transfected with human FSH R. Morgenthaler, N.G. <i>et al.</i> (1998) <i>Horm. Metab. Res.</i> 30 :162. The Neutralization Dose (ND ₅₀) is typically 0.02-0.1 µg/mL in the presence of 1 ng/mL Recombinant Human FSH α/β.
-----------------------	---

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Follicle stimulating hormone receptor (FSH R), also called follitropin receptor or LGR1, is a 695 amino acid (aa), ~87 kDa 7-transmembrane receptor of the leucine-rich repeat glycoprotein hormone receptor subfamily of GPCRs. It is expressed by ovarian granulosa cells and is essential for ovarian follicle maturation. In the male, it is expressed by Sertoli cells and plays a minor role in male fertility. Human FSH R shares 86% aa sequence identity with mouse and rat FSH R within the 349 aa N-terminal extracellular domain. A 754 aa isoform contains an inserted sequence at aa 75.