

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CD79A in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant mouse CD79A is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 706931
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CD79A Leu33-Arg143 Accession # P11912
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CD79A, also known as Igα and MB-1, is a 44 kDa transmembrane glycoprotein in the immunoglobulin superfamily. It contains a single Ig-like domain in its extracellular region (ECD) and one cytoplasmic immunoreceptor tyrosine-based activation motif (ITAM). Alternate splicing generates a short isoform with a 39 aa deletion in the ECD. Heterodimers of CD79A and CD79B/Igβ associate with a membrane bound immunoglobulin on the B cell surface to form the B cell antigen receptor complex (BCR). CD79A and CD79B are required for BCR-mediated signaling and consequently for the development and activation of B lineage cells. Within the ECD, human CD79A shares 57% aa sequence identity with mouse and rat CD79A.