

DESCRIPTION

Species Reactivity	Porcine
Specificity	Detects porcine IL-10 in direct ELISAs. In direct ELISAs, 100% cross-reactivity with recombinant canine IL-10, recombinant feline IL-10, recombinant human IL-10, and recombinant viral IL-10 is observed and no cross-reactivity with recombinant cotton rat IL-10, recombinant equine IL-10, recombinant mouse IL-10, recombinant rat IL-10, or recombinant human IL-22 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 262715
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant porcine IL-10 Ser19-Asn175 Accession # Q29055
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.


	Recommended Concentration	Sample
Immunocytochemistry	8-25 µg/mL	See Below

DATA

Immunocytochemistry



Treated



Untreated (control)

IL-10 in Porcine PBMCs.
IL-10 was detected in immersion fixed porcine peripheral blood mononuclear cells treated with calcium ionomycin and PMA using Mouse Anti-Porcine IL-10 Monoclonal Antibody (Catalog # MAB6932) at 15 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). Specific staining was localized to cytoplasm. View our protocol for [Fluorescent ICC Staining of Non-adherent Cells](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

IL-10, initially designated cytokine synthesis inhibitory factor (CSIF), was originally identified as a product of murine T helper 2 (Th2) clones that inhibited the cytokine production by Th1 clones which are dependent upon stimulation with antigen in the presence of antigen presenting cells (APC). Murine IL-10 is produced by Th2 cells, activated fetal thymocytes, macrophages, keratinocytes, and LY-1⁺ (CD5⁺) and normal B cells. Human IL-10 has cross-species activities and is active on mouse cells. Murine IL-10 is species-specific and does not act on human cells. Porcine IL-10 shares 71% and 78% amino acid sequence identity with mouse and human IL-10, respectively. IL-10 is a pleiotropic cytokine that can exert either immunostimulatory or immunosuppressive effects on a variety of cell types. It is a potent immunosuppressant of macrophage functions. *In vitro*, IL-10 can inhibit the accessory function and antigen-presenting capacity of monocytes by, among other effects, down-regulating class II MHC expression. Thus, IL-10 can inhibit monocyte/macrophage-dependent, antigen stimulated cytokine synthesis (especially IFN-γ) by human PBMNC and NK, and mouse Th1 cells. Additionally, IL-10 is a potent inhibitor of monocyte/macrophage activation and its resultant cytotoxic effects. As an immunostimulatory cytokine, IL-10 can act on B cells to enhance their viability, cell proliferation, Ig secretion, and class II MHC expression. Aside from B lymphocytes, IL-10 is also a growth co-stimulator for thymocytes and mast cells, as well as an enhancer of cytotoxic T cell development.

References:

1. Moore, K.W. *et al.* (1993) *Annu. Rev. Immunol.* **11**:165.