

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human ICOS in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 669222
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with human ICOS and mouse myeloma cell line NS0-derived recombinant human ICOS cocktail Glu21-Phe141 Accession # Q9Y6W8
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Inducible costimulator (ICOS), also called AILIM (activation-inducible lymphocyte immunomediatory molecule) and CRP-1 (CD28-related protein-1), is a member of the growing CD28 family of immune costimulatory receptors. Other family members are CD28, CTLA-4, and PD-1. Human ICOS is a homodimeric type I transmembrane protein consisting of 199 amino acids (aa) with a putative 20 aa signal sequence, a 121 aa extracellular domain, a 23 aa transmembrane region, and a 35 aa cytoplasmic domain. ICOS shares approximately 39% amino acid similarity with CD28 and CTLA-4. Human and mouse ICOS share approximately 72% amino acid identity. ICOS is expressed on most CD45RO⁺ cells. ICOS expression is up-regulated within approximately 24-48 hours of activation on T_H primed cells. B7-H2, a member of the B7 family of costimulatory ligands, has been identified as the ICOS ligand. The B7-H2/ICOS interaction appears to play roles in T cell dependent B cell activation and T_H differentiation.

References:

1. Aicher, A. *et al.* (2000) J. Immunol. **164**:4689.
2. Coyle, A.J. *et al.* (2000) Immunity **13**:95.
3. Coyle, A.J. and J.C. Gutierrez-Ramos (2001) Nat. Immunol. **2**:203.
4. Gonzalo, J.A. *et al.* (2001) J. Immunol. **166**:1.
5. Hutloff, A. *et al.* (1999) Nature **397**:263.
6. Mages, H.W. *et al.* (2000) Eur. J. Immunol. **30**:1040.
7. Yoshinaga, S.K. *et al.* (1999) Nature **402**:827.