

**DESCRIPTION**

|                           |   |
|---------------------------|---|
| <b>Species Reactivity</b> | Human   |
| <b>Specificity</b>        | Detects human STIM1 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human STIM2 is observed.   |
| <b>Source</b>             | Monoclonal Mouse IgG <sub>2B</sub> Clone # 705129   |
| <b>Purification</b>       | Protein A or G purified from hybridoma culture supernatant  |
| <b>Immunogen</b>          | <i>E. coli</i> -derived recombinant human STIM1<br>Leu23-Thr182<br>Accession # Q13586   |
| <b>Formulation</b>        | Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.<br>*Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS. |


**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

|                             | <b>Recommended Concentration</b> | <b>Sample</b> |
|-----------------------------|----------------------------------|---------------|
| <b>Immunohistochemistry</b> | 8-25 µg/mL                       | See Below     |

**DATA**

**Immunohistochemistry**



**STIM1 in Human Brain.** STIM1 was detected in immersion fixed paraffin-embedded sections of human brain (cerebellum) using Mouse Anti-Human STIM1 Monoclonal Antibody (Catalog # MAB69761) at 15 µg/mL overnight at 4 °C. Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Specific staining was localized to cytoplasm in Purkinje neurons. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

**PREPARATION AND STORAGE**

|                                |  |
|--------------------------------|--|
| <b>Reconstitution</b>          | Sterile PBS to a final concentration of 0.5 mg/mL.   |
| <b>Shipping</b>                | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.<br>*Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C   |
| <b>Stability &amp; Storage</b> | <b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul> |

**BACKGROUND**

STIM-1 (stromal interaction molecule 1), previously called GOK, is a 90 kDa type I transmembrane protein of the endoplasmic reticulum (ER). When STIM-1 senses depletion of calcium in the ER via its EF-hand domain (aa 63-98), it interacts with the plasma membrane Ca<sup>2+</sup> release-activated Ca<sup>2+</sup> (CRAC) channel Orai1, increasing Ca<sup>2+</sup> influx. The human STIM-1 extracellular/luminal domain (aa 23-213) contains an EF hand Ca<sup>2+</sup>-binding motif and a SAM (sterile a-motif) multimerization domain. A potential isoform is truncated at aa 491. Defects in STIM1 are the cause of immune dysfunction with T-cell inactivation due to calcium entry defect type 2 (IDTICED2). Within the region used as an immunogen, human STIM1 shares 95% amino acid identity with mouse and rat STIM1.