

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human KLRG1 in direct ELISAs and Western blots.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 697213
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human KLRG1 Leu60-Phe195 Accession # Q96E93
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	2 µg/mL	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

KLRG1 (killer cell lectin-like receptor G1), also known as MAFA (mast cell function associated), is a 30-38 kDa type II transmembrane inhibitory glycoprotein of the C-type lectin family, designated CLEC15A. KLRG1 cDNA encodes 195 amino acids (aa) including an intracellular ITIM motif and a 136 aa extracellular domain (ECD) with a single C-type lectin domain. The human KLRG1 ECD shares 57% and 54% aa identity with mouse and rat KLRG1, respectively. A 189 aa isoform diverges at aa 186. KLRG1 binds E-, N- and R-cadherins and functions as an MHC-independent means of identifying non-self pathogens and epithelial tumor cells with low E-cadherin expression. It is expressed as a monomer or disulfide-linked homodimer on NK and T cell subsets such as tumor-infiltrating lymphocytes.