

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human ABCA1 in direct ELISAs.
<b>Source</b>	Monoclonal Mouse IgG <sub>2A</sub> Clone # 768510
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human ABCA1 Phe1248-Gln1350 Accession # O95477
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Intracellular Staining by Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.5 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

ABCA1 (ATP-binding cassette, subfamily A member 1; also cholesterol efflux regulatory protein) is a 240-280 kDa member of the ABCA family, ABC transporter superfamily of molecules. It is widely expressed, particularly by macrophages, where it participates in the transport of lipid across cell membranes. Once transported, it is passed on to tethered apoA-I to form HDL. Human ABCA1 is 2261 amino acids (aa) in length. It has up to 15 transmembrane segments with beginning and ending cytoplasmic domains and a large cytoplasmic central region. ABC transporters are found between aa 899-1311 and 1912-2144. Amino acids 2216-2221 are essential for lipid transport, and two intramolecular disulfide bonds are required for activity (Cys75-Cys309 and Cys1463/1465-Cys1477). There are three potential isoform variants. One possesses an alternative start site at Met61, while two others contain 12 aa and 123 aa substitutions for aa 241-2261, respectively. Over aa 1248-1350, human ABCA1 shares 97% aa identity with mouse ABCA1.