

#### DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human ABCA1 in direct ELISAs and Western blots.
<b>Source</b>	Recombinant Monoclonal Rabbit IgG Clone # 1276B
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	Human ABCA1 synthetic peptide Accession # O95477
<b>Formulation</b>	Supplied as a solution in PBS containing BSA, Glycerol and Sodium Azide. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	See Below

#### DATA



#### PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 °C, as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after opening.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after opening.</li> </ul>

#### BACKGROUND

ABCA1 (ATP-binding cassette, subfamily A member 1; also cholesterol efflux regulatory protein) is a 240-280 kDa member of the ABCA family, ABC transporter superfamily of molecules. It is widely expressed, particularly by macrophages, where it participates in the transport of lipid across cell membranes. Once transported, it is passed on to tethered apoA-I to form HDL. Human ABCA1 is 2261 amino acids (aa) in length. It has up to 15 transmembrane segments with beginning and ending cytoplasmic domains and a large cytoplasmic central region. ABC transporters are found between aa 899-1311 and 1912-2144. Amino acids 2216-2221 are essential for lipid transport, and two intramolecular disulfide bonds are required for activity (Cys75-Cys309 and Cys1463/1465-Cys1477). There are three potential isoform variants. One possesses an alternative start site at Met61, while two others contain 12 aa and 123 aa substitutions for aa 241-2261, respectively. Over aa 1248-1350, human ABCA1 shares 97% aa identity with mouse ABCA1.

#### PRODUCT SPECIFIC NOTICES

\* Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to SDS for additional information and handling instructions.