

DESCRIPTION



Species Reactivity	Mouse
Specificity	Detects mouse Kallikrein 5 in ELISA.
Source	Monoclonal Rat IgG _{2B} Clone # 880036
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Kallikrein 5 23-293
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	2 µg/mL	See Below
Immunohistochemistry	8-25 µg/mL	See Below

DATA

<p>Western Blot</p> 	<p>Detection of Mouse Kallikrein 5 by Western Blot. Western blot shows lysates of NMuMG mouse mammary gland epithelial cell line. PVDF membrane was probed with 2 µg/mL of Rat Anti-Mouse Kallikrein 5 Monoclonal Antibody (Catalog # MAB7236) followed by HRP-conjugated Anti-Rat IgG Secondary Antibody (Catalog # HAF005). A specific band was detected for Kallikrein 5 at approximately 30 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p>Immunohistochemistry</p>  <p>Kallikrein 5 in Hair Follicle of Mouse Skin. Kallikrein 5 was detected in immersion fixed frozen sections of mouse skin using Rat Anti-Mouse Kallikrein 5 Monoclonal Antibody (Catalog # MAB7236) at 25 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Rat HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS017) and counterstained with hematoxylin (blue). Specific staining was localized to the inner sheath cuticle of hair follicles. View our protocol for Chromogenic IHC Staining of Frozen Tissue Sections.</p>
---	---	---

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Tissue Kallikrein 5 (KLK5) is a secreted serine protease with trypsin-like substrate specificity. KLK5 is highly expressed in the skin, and is also known as stratum corneum tryptic enzyme due to its high expression level in this tissue. KLK5 is thought to play an important role in the process of desquamation (1). KLK5 is a potential biomarker for ovarian and breast cancers (2). It may also be a physiological activator of KLK7 in the stratum corneum (3). Recombinant mouse KLK5 is purified as the proform, and can undergo autoactivation under the appropriate conditions. KLK5 enzymatic activity can be inhibited by the following recombinant human serpins: Serpin-A1, -A5, -F2, and -G1 (Catalog # 1268-PI, 1266-PI, 1470-PI, and 2488-PI).

References:

1. Brattsand M. and T. Egelrud (1999) J. Biol. Chem. **274**:30033.
2. Yousef G.M. *et al.* (2003) Cancer Res. **63**:3958.
3. Kim H. *et al.* (2001) Br. J. Cancer **84**:543.