

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Attractin in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 750003
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Attractin Ala84-Gln1272 Accession # O75882
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Immunocytochemistry	8-25 µg/mL	See Below

DATA

Immunocytochemistry



Attractin in HepG2 Human Cell Line.
Attractin was detected in immersion fixed HepG2 human hepatocellular carcinoma cell line using Mouse Anti-Human Attractin Monoclonal Antibody (Catalog # MAB7238) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the Northern-Lights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). View our protocol for [Fluorescent ICC Staining of Cells on Coverslips](#).

PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Attractin (ATRN), also known as DPPT-L, is an approximately 200 kDa transmembrane glycoprotein that shows dipeptidyl peptidase activity similar to DPPIV/CD26. Attractin is involved in a variety of processes including monocyte-T cell adhesion, axon myelination, melanocyte pigment synthesis, and energy homeostasis. The extracellular region of human Attractin contains one EGF-like domain, one CUB domain, six Kelch repeats, four PSI domains, one C-type lectin domain, and two laminin EGF-like domains. Alternate splicing of human Attractin generates a secreted isoform that lacks the transmembrane and cytoplasmic regions. Attractin is transiently upregulated during T cell activation before expression switches to the 175 kDa secreted isoform which is released into the circulation. Soluble Attractin is preferentially expressed by leukocytes and differentiating neurons. It blocks neurite formation and is elevated in the CSF of astrocytoma patients. Within aa 84-1272, human Attractin shares 95% aa sequence identity with mouse and rat Attractin.