

**DESCRIPTION**



<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse TGF- $\beta$ 2 in ELISAs and Western blots. In direct ELISAs, 100% cross-reactivity with recombinant human (rh) TGF- $\beta$ 2, 25% cross-reactivity with rhTGF- $\beta$ 3, and no cross-reactivity with recombinant mouse TGF- $\beta$ 1 is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2B</sub> Clone # 771244
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant mouse TGF- $\beta$ 2 Ala303-Ser414 Accession # P27090
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 $\mu$ m filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	2 $\mu$ g/mL	See Below
<b>Immunohistochemistry</b>	8-25 $\mu$ g/mL	See Below

**DATA**

<p><b>Western Blot</b></p> 	<p><b>Detection of Human and Mouse TGF-<math>\beta</math>2 by Western Blot.</b> Western blot shows lysates of Y-79 human retinoblastoma cell line, HUVEC human umbilical vein endothelial cells, NIH-3T3 mouse embryonic fibroblast cell line, 4T1 mouse breast cancer cell line, and NMuMG mouse mammary gland epithelial cell line. PVDF membrane was probed with 2 <math>\mu</math>g/mL of Rat Anti-Mouse TGF-<math>\beta</math>2 Monoclonal Antibody (Catalog # MAB73461) followed by HRP-conjugated Anti-Rat IgG Secondary Antibody (Catalog # HAF005). A specific band was detected for TGF-<math>\beta</math>2 at approximately 52 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p><b>Immunohistochemistry</b></p>  <p><b>TGF-<math>\beta</math>2 in Mouse Embryo.</b> TGF-<math>\beta</math>2 was detected in immersion fixed frozen sections of mouse embryo (15 d.p.c.) using Rat Anti-Mouse TGF-<math>\beta</math>2 Monoclonal Antibody (Catalog # MAB73461) at 25 <math>\mu</math>g/mL overnight at 4 °C. Tissue was stained using the Anti-Rat HRP-DAB Cell &amp; Tissue Staining Kit (brown; Catalog # CTS017) and counterstained with hematoxylin (blue). Specific staining was localized to neuronal processes. View our protocol for <a href="#">Chromogenic IHC Staining of Frozen Tissue Sections</a>.</p>
---	--	---

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.5 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

TGF- $\beta$ 2 (transforming growth factor beta 2) is one of three closely related mammalian members of the large TGF- $\beta$  superfamily that share a characteristic cysteine knot structure. TGF- $\beta$ 1, -2 and -3 are highly pleiotropic cytokines proposed to act as cellular switches that regulate processes such as immune function, proliferation and epithelial-mesenchymal transition. Each TGF- $\beta$  isoform has some non-redundant functions; for TGF- $\beta$ 2, mice with targeted deletion show defects in development of cardiac, lung, craniofacial, limb, eye, ear and urogenital systems. Mouse TGF- $\beta$ 2 cDNA encodes a 414 amino acid (aa) precursor that contains a 19 aa signal peptide and a 395 aa proprotein. A furin-like convertase processes the proprotein to generate an N-terminal 283 aa latency-associated peptide (LAP) and a C-terminal 112 aa mature TGF- $\beta$ 2. Disulfide-linked homodimers of LAP and TGF- $\beta$ 2 remain non-covalently associated after secretion, forming the small latent TGF- $\beta$ 2 complex. Covalent linkage of LAP to one of three latent TGF- $\beta$  binding proteins (LTBPs) creates a large latent complex that may interact with the extracellular matrix. TGF- $\beta$  is activated from latency by pathways that include actions of the protease plasmin, matrix metalloproteases, thrombospondin 1 and a subset of integrins. Mature mouse TGF- $\beta$ 2 shares 100% aa identity with rat TGF- $\beta$ 2, and 97% aa identity with human, porcine, canine, equine and bovine TGF- $\beta$ 2. It demonstrates cross-species activity. In most cells, TGF- $\beta$ 2 signaling begins with binding to a complex of the accessory receptor betaglycan (also known as TGF- $\beta$  RIII) and a type II ser/thr kinase receptor termed TGF- $\beta$  RII, which then phosphorylates and activates another ser/thr kinase receptor, TGF- $\beta$  RI (also called activin receptor-like kinase (ALK) -5), or alternatively, ALK-1. The whole complex phosphorylates and activates Smad proteins that regulate transcription. In bone-related cells, however, TGF- $\beta$ 2 also signals through TGF- $\beta$  RIIB (a splice variant of TGF- $\beta$  RII), independently of TGF- $\beta$  RIII. Use of other signaling pathways that are Smad-independent allows for disparate actions observed in response to TGF- $\beta$  in different contexts.