

DESCRIPTION


Species Reactivity	Canine
Specificity	Detects canine IL-10 in Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 138128
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant canine IL-10 Ser20-Ile179 Accession # XP_855560
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	Recombinant Canine IL-10 (Catalog # 735-CL)
Neutralization		Measured by its ability to neutralize IL-10-induced proliferation in the MC/9-2 mouse mast cell line. Thompson-Snipes, L. <i>et al.</i> (1991) <i>J. Exp. Med.</i> 173 :507. The Neutralization Dose (ND ₅₀) is typically 0.1-0.25 µg/mL in the presence of 2 ng/mL Recombinant Canine IL-10.

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

IL-10, initially designated cytokine synthesis inhibitory factor (CSIF), was originally identified as a product of murine T helper 2 (Th2) clones that inhibited the cytokine production by Th1 clones. The canine homologue of mouse IL-10 has been cloned. Canine IL-10 shares 80% amino acid sequence homology with human IL-10 and 72% sequence homology with mouse IL-10.

IL-10 is a pleiotropic cytokine that can exert either immunostimulatory or immunosuppressive effects on a variety of cell types. It is a potent immunosuppressant of macrophage functions. *In vitro*, IL-10 can inhibit the accessory function and antigen-presenting capacity of monocytes by, among other effects, downregulating class II MHC expression. Thus, IL-10 can inhibit monocyte/macrophage-dependent, antigen stimulated cytokine synthesis (especially IFN-γ) by human PBMC and NK, and mouse Th1 cells. Additionally, IL-10 is a potent inhibitor of monocyte/macrophage activation and its resultant cytotoxic effects. As an immunostimulatory cytokine, IL-10 can act on B cells to enhance their viability, cell proliferation, Ig secretion, and class II MHC expression. Aside from B lymphocytes, IL-10 is also a growth co-stimulator for thymocytes and mast cells, as well as an enhancer of cytotoxic T cell development.

References:

1. Moore, K.W. *et al.* (1993) *Annu. Rev. Immunol.* **11**:165.