

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse GDF-9 in direct ELISAs.
Source	Monoclonal Rat IgG _{2A} Clone # 785123
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse GDF-9 Met1-Arg441 Accession # Q07105
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Neutralization	Measured by its ability to neutralize GDF-9-induced growth inhibition in the Mv1Lu mink lung epithelial cell line. The Neutralization Dose (ND ₅₀) is typically 0.4-2.4 µg/mL in the presence of 0.5 µg/mL Recombinant Mouse GDF-9.
-----------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Growth differentiation factor-9 (GDF-9) is a member of the transforming growth factor- β (TGF- β) superfamily, and is an oocyte secreted paracrine factor essential for mammalian ovarian folliculogenesis (1-2). Mouse GDF-9 is synthesized as a 441 amino acid (aa) prepropeptide that contains a 29 aa signal sequence, a 277 aa propeptide, and a 135 aa mature chain. Residues 340-441 constitute a TGF- β like domain. In addition, there is one potential site of N-linked glycosylation in the mature chain. Unlike other members of the TGF- β superfamily, GDF-9 lacks the conserved cysteine residue that is believed to form the sole disulfide linkage between subunits in other family members (3). Mature mouse GDF-9 shares 90% aa sequence identity with mature human GDF 9. The protein is expressed throughout the development of the maturing follicle (2). GDF-9 functions as a paracrine factor in the regulation of granulosa cell proliferation and differentiation, and is essential for fertility (2, 4). Studies on GDF-9 null mice have demonstrated arrested follicular development at the primary follicle stage (5). Mouse GDF-9 induces Smad2 phosphorylation and inhibin production in rat diethylstilbestrol treated granulosa cells (6) and in human granulosa-luteal cells (7). The downstream signaling actions of GDF 9 are mediated by the type I receptor, activin receptor-like kinase 5 (ALK5), initiating the subsequent activation of Smad2 and Smad3 (2, 8). GDF 9 uses the BMP type II receptor (BMPRII) as its other signaling receptor (2, 9).

References:

1. McGrath, S.A. *et al.* (1995) *Mol. Endocrinol.* **9**:131.
2. Mottershead, D.G. *et al.* (2008) *Mol. Cell. Endocrinol.* **283**:58.
3. McPherron, A.C. and S.-J. Lee (1992) *J. Biol. Chem.* **268**:3444.
4. Gilchrist, R.B. *et al.* (2006) *J. Cell. Sci.* **119**:3811.
5. Dong, J. *et al.* (1996) *Nature* **383**:531.
6. Roh, J.S. *et al.* (2003) *Endocrinology* **144**:172.
7. Kaivo-Oja, N. *et al.* (2003) *J. Clin. Endocrinol. Metab.* **88**:755.
8. Mazerbourg, S. *et al.* (2004) *Mol. Endocrinol.* **18**:653.
9. Vitt, U.A. *et al.* (2002) *Biol. Reprod.* **67**:473.