

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human BATF3 in ELISA.
Source	Monoclonal Mouse IgG _{2B} Clone # 841702
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human BATF3 Met1-Arg127 Accession # Q9NR55
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

BATF3 (Basic leucine zipper transcriptional factor ATF-like 3; also p21SNFT) is a 20 kDa nuclear member of the bZIP family of proteins. It is expressed in Th1 cells and conventional dendritic cells (CD11c⁺), and serves to downregulate AP-1 mediated transcription. BATF3 accomplishes this by heterodimerizing with Jun and binding to AP-1 consensus binding sites, thus precluding a Jun/Fos interaction with gene activation. Human BAFT3 is 127 amino acids (aa) in length. It contains one DNA binding motif (aa 41-59) with an adjacent leucine-zipper (aa 63-84), but lacks a transactivation domain. Full-length human BATF3 (aa 1-127) shares 80% aa sequence identity with mouse BATF3.