

DESCRIPTION



Species Reactivity	Human
Specificity	Detects human RIPK3/RIP3 in direct ELISAs.
Source	Recombinant Monoclonal Rabbit IgG Clone # 1312D
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Synthetic peptide containing human RIPK3/RIP3 Accession # Q9Y572
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
Western Blot	2 µg/mL	See Below
Immunocytochemistry	1-25 µg/mL	See Below

DATA

<p>Western Blot</p>  <p>Detection of Human RIPK3/RIP3 by Western Blot. Western blot shows lysates of SK-BR-3 human breast cancer cell line and Raji human Burkitt's lymphoma cell line. PVDF membrane was probed with 2 µg/mL of Rabbit Anti-Human RIPK3/RIP3 Monoclonal Antibody (Catalog # MAB76042) followed by HRP-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog # HAF008). A specific band was detected for RIPK3/RIP3 at approximately 60 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p>Immunocytochemistry</p>  <p>RIPK3/RIP3 in SK-BR-3 and HepG2 Human Cell Lines. RIPK3/RIP3 was detected in immersion fixed SK-BR-3 human breast cancer cell line (positive stain) and HepG2 human hepatocellular carcinoma cell line (negative stain) using Rabbit Anti-Human RIPK3/RIP3 Monoclonal Antibody (Catalog # MAB76042) at 1 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Rabbit IgG Secondary Antibody (red; Catalog # NL004) and counterstained with DAPI (blue). Specific staining was localized to cytoplasm. View our protocol for Fluorescent ICC Staining of Cells on Coverslips.</p>
---	---

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

RIP3 (Receptor-Interacting Protein 3), also known as RIPK3, is a 518 amino acid (aa), 60 kDa phosphoprotein that contains an N-terminal protein kinase domain and a C-terminal interaction motif with which it binds RIP1. It converts cell response to TNF-α from apoptosis (RIP1 via Fas) to programmed necrosis (RIP1/RIP3) by inducing reactive oxygen species production. Human RIP3 (aa 1-218) shares approximately 74% aa sequence identity with mouse and rat RIP3. Alternately spliced human b (252 aa) and g (231 aa) isoforms diverge after aa 219 and 221, respectively, and are found to antagonize RIP3.