

**DESCRIPTION**


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse CD161/NK1.1 expressed specifically in the C57/B6 mouse strain. No detection of CD161/NK1.1 was observed in the BALB/c mouse strain. In direct ELISAs, less than 5% cross-reactivity with recombinant mouse (rm) KLRB-1B is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 694370
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse CD161/NK1.1 Val62-Ser220 Accession # P27814
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.5 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

KLRB1C, also known as NK1.1, CD161c, Ly-55, and NKR-P1c, is a 28 kDa type 2 transmembrane protein in the killer cell lectin-like receptor family. CD161 is expressed as a disulfide-linked homodimer on the surface of NK cells and subpopulations of NKT, CD4<sup>+</sup>, CD8<sup>+</sup>, and γδ T cells. Its cross-linking on NK cells induces cytolytic activity, and on CD4 cells it promotes IgE production and the expansion of Th2 responses. Mature mouse CD161 consists of a 45 amino acid (aa) cytoplasmic domain, a 21 aa transmembrane segment, and a 157 aa extracellular domain with one C-type lectin domain.