

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human Connexin 43/GJA1. Stains human Connexin 43/GJA1 transfectants but not irrelevant transfectants in flow cytometry.
Source	Monoclonal Mouse IgG _{2A} Clone # 578618
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0 transfected with human Connexin 43/GJA1 Accession # P17302
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Connexin43 (Cx43; also gap junction α -1 protein) is a 41 - 44 kDa member of the connexin family, α -type subfamily, of transmembrane proteins. It is the most common type of connexin in cardiac muscle cells, and also occurs in hepatocytes, astrocytes and ovary granulosa cells. It is a 4-transmembrane protein 382 amino acids (aa) in length that contains two cytoplasmic tails. One is 12 aa in length at the N-terminus and the second is 151 aa in length at the C-terminus. Human and rat Cx43 are 98% aa identical over the entire length of the molecule. Connexins form gap junctions (GJs) which are intercellular channels between cells. Each adjacent cell contributes to a functional channel. The fundamental unit is a "connexon", or hemi-channel, which is composed of six connexins in a sliding subunit hexamer configuration. The hexamer may be either homomeric or heteromeric. The connexon arrangement provides for the opening and closing of an intersubunit space (or pore) that allows diffusion of molecules 1 kDa or less. It is suggested that the extended C-terminus of each connexin may interact with multiple docking proteins and serve as a plug during closure. Connexons on adjacent cells interact via their extracellular loops to form a GJ channel. When open, these channels allow for the transit of small hydrophilic molecules such as ATP, glucose, IP3 and Ca⁺⁺. Connexons (or hemi-channels) that do not interact with adjacent cells are also suggested to allow for small molecule transit under unusual circumstances. Cx43 activity is regulated by phosphorylation. The exact effects, however, may be context-specific. In one case, PKC phosphorylation of S368 has been found to decrease GJ channel permeability to select small molecular weight solutes. Alternatively, PKA has been suggested to be the mediator of Cx43 phosphorylation of S365, S368, S369 and S373. In this case, phosphorylation promotes channel activity.

References:

1. Sohl, G. and K. Willecke (2004) *Cardiovasc. Res.* **62**:228.
2. Willecke, K. *et al.* (2002) *Biol. Chem.* **383**:725.
3. Giepmans, B.N.G. (2004) *Cardiovasc. Res.* **62**:233.
4. Vinken, M. *et al.* (2006) *Cell. Signal.* **18**:592.
5. Stout, C. *et al.* (2004) *Curr. Opin. Cell Biol.* **16**:507.
6. Bao, X. *et al.* (2004) *Am. J. Physiol. Cell. Physiol.* **286**:C647.
7. Yogo, K. *et al.* (2006) *J. Reprod. Dev.* **52**:321.