

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human ALK-7 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant rat ALK-7 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 810506
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human ALK-7 Leu26-Glu113 Accession # Q8NER5
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.5 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

Activin receptor-like kinase 7 (ALK-7), also known as Activin R1C (gene name ACVR1C), is a glycosylated 58 kDa type I receptor in the superfamily of TGF-β serine/threonine kinase receptors. It associates with type II receptors to form a signaling complex that responds to the ligands Activin AB, and Activin B, GDF3, and Nodal. ALK-7 plays a role in regulating energy balance by inhibiting insulin secretion and inducing pancreatic beta cell apoptosis. It is expressed in adipose tissue but downregulated in obesity. ALK-7 is also expressed in pituitary gonadotropic cells and in pre-eclamptic placenta. It induces the apoptosis of trophoblasts as well as ovarian granulosa and epithelial cells. Within the extracellular domain, human ALK-7 shares 95% and 91% amino acid (aa) sequence identity with mouse and rat ALK-7, respectively. Alternate splicing of human ALK-7 generates additional isoforms with either a 50 aa N-terminal truncation or with deletions of 79 aa or 157 aa that encompass the transmembrane segment.