

## DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Doc2α in ELISAs. In direct ELISAs, no cross-reactivity with recombinant human Doc2β is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 842213
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human Doc2α Met1-Lys114 Accession # Q14183
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 μm filtered solution in PBS.


## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Immunohistochemistry</b>	8-25 μg/mL	See Below

## DATA

### Immunohistochemistry



**Doc2α in Human Brain.** Doc2α was detected in immersion fixed paraffin-embedded sections of human brain (hippocampus) using Mouse Anti-Human Doc2α Monoclonal Antibody (Catalog # MAB7904) at 15 μg/mL overnight at 4 °C. Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Specific staining was localized to cytoplasm of neurons. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

## PREPARATION AND STORAGE

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.5 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

DOC2A/Doc2α (Double C2-like domain containing protein alpha) is a 44 kDa (predicted) monomeric member of the C2 domain-containing protein family of molecules. It is expressed in both neurons and mast cells and appears to serve as an intracellular Ca<sup>++</sup> sensor protein that regulates secretory vesicle release. In neurons, Doc2α is normally bound to synaptic vesicles and interacts with Munc13-1 to promote secretory vesicle exocytosis through the cell membrane. In mast cells, a similar process occurs that involves Munc13-4 instead of Munc13-1. Human Doc2α is 400 amino acids (aa) in length. It contains a Mid domain (aa 13-37) that binds Munc13-1, followed by one C2 domain that binds Ca<sup>++</sup> and lipid (aa 91-195) and a second C2 domain that binds SNAP25 (253-356). There is one potential alternative start site 16 aa upstream of the standard site. Over aa 1-114, human Doc2α shares 90% aa sequence identity with mouse Doc2α. Human DOC2B is the product of a separate gene and shares no meaningful aa sequence identity (<30%) with human Doc2α.