

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Insulin in ELISA.
Source	Monoclonal Mouse IgG ₁ Clone # 873724
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	The construct consists of the alpha mating factor signal followed by a 10 aa peptide leader, B chain aa 1-29, AAK spacer A chain aa 1-21.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	8-25 µg/mL	See Below

DATA

Immunohistochemistry



Insulin in Human Pancreas. Insulin was detected in immersion fixed paraffin-embedded sections of human pancreas using Mouse Anti-Human Insulin Monoclonal Antibody (Catalog # MAB8056) at 15 µg/mL overnight at 4 °C. Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counter-stained with hematoxylin (blue). Specific staining was localized to islets. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Insulin is a disulfide-linked heterodimeric protein secreted by the pancreatic Islets of Langerhans. Mature insulin is generated by the proteolytic removal of a peptide from proinsulin. It is involved in the regulation of glucose metabolism through interactions with the Insulin Receptor.