

DESCRIPTION


| | |
|---------------------------|---|
| Species Reactivity | Rat |
| Specificity | Detects rat CXCR3 in ELISA. Stains rat CXCR3 transfected cells but not irrelevant transfectants by Flow Cytometry. |
| Source | Monoclonal Mouse IgG _{2B} Clone # 868013 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | NS0 mouse myeloma cell line transfected with rat CXCR3 Accession # Q9J119 |
| Formulation | Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|--|---------------|
| Flow Cytometry | 2.5 µg/10 ⁶ cells | See Below |
| CytoF-ready | Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation. | |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Reconstitution | Reconstitute at 0.5 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

CXCR3, also known as CD183, is an approximately 41 kDa (predicted) 7-transmembrane spanning receptor for the chemokines CXCL9, CXCL10, and CXCL11. It is expressed on activated T cells, B cells, and NK cells during inflammation. It is additionally upregulated on solid tumor cells, tumor endothelium, and cancer stem cells. CXCR3 plays a role in leukocyte recruitment to sites of inflammation. It also contributes to Th1 biased responses during autoimmune diseases rheumatoid arthritis, systemic erythematosis, and type 1 diabetes. Rat CXCR3 shares 86% and 96% amino acid sequence identity with human and mouse CXCR3, respectively.