

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human cIAP-1/HIAP-2 in direct ELISAs. In direct ELISAs, 25% cross-reactivity with recombinant human cIAP-2 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 681724
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human cIAP-1/HIAP-2 His2-Ser618 Accession # Q13490
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	8-25 µg/mL	See Below

DATA

Immunohistochemistry



cIAP-1/HIAP-2 in Human Prostate. cIAP-1/HIAP-2 was detected in immersion fixed paraffin-embedded sections of human prostate using Mouse Anti-Human cIAP-1/HIAP-2 Monoclonal Antibody (Catalog # MAB8181) at 15 µg/mL overnight at 4 °C. Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Specific staining was localized to the cytoplasm of epithelial cells. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

cIAP-1 (also known as BIR2, MIHB and HIAP-2) is a member of the inhibitor of apoptosis (IAP) family of proteins that inhibit the proteolytic activity of mature caspases. cIAP-1 has 3 BIR (baculovirus inhibitor of apoptosis) domains, a RING finger domain, and a caspase recruitment domain (CARD). cIAP-1 inhibits caspases by interaction of the BIR domain with the active caspase. Caspase activity may be restored through interactions with the Reaper like motif on mitochondrial proteins such as SMAC/Diablo or HTRA-2/Omi. cIAP-1 is reported to be cleaved by caspases in fetal rat hepatocytes treated with TGF-β.

References:

- Roy, N. *et al.* (1997) EMBO J. **23**:6914.
- Deveraux, Q. *et al.* (1997) Nature **388**:300.
- Deveraux, Q. and J. Reed (1999) Genes & Develop. **13**:239.
- Herrera, B. *et al.* (2002) FEBS Letters **520**:93.