

DESCRIPTION


Species Reactivity	Human
Specificity	Stains human Melanocortin-5 R/MC5R transfected cells but not irrelevant transfectants in flow cytometry.
Source	Monoclonal Mouse IgG _{2A} Clone # 821012
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with human Melanocortin-5 R/MC5R Met1-Asp325 Accession # P33032
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Melanocortin 5R (MC5R) is a member of the melanocortin receptor family. The 325 amino acid G-protein-coupled 7-transmembrane glycoprotein is expressed in skeletal muscle, where engagement by α-melanocyte stimulating hormone increases glucose uptake and stimulates a thermogenic response. Expression is also found in sebaceous glands and other skin structures, and on selected hematopoietic cell lines. Human MC5R shares 80-81% amino acid sequence identity with mouse and rat MC5R.