

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human $\beta_2$ -Microglobulin in direct ELISAs and Western blots.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 883028
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human $\beta_2$ -Microglobulin Met1-Met119 Accession # P61769
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 $\mu$ m filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	2 $\mu$ g/mL	See Below
<b>Flow Cytometry</b>	2.5 $\mu$ g/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

$\beta_2$ -Microglobulin (b2M) is a 12 kDa secreted polypeptide that serves as the light chain of Class I MHC molecules. Possessing an Ig-like domain, b2M noncovalently associates with both 44 kDa classical (HLA-A, -B, -C) and 40 kDa non-classical (HLA-E, -F, -G) Class I MHC heavy chains as well as with 43-49 kDa Class I non-MHC heavy chains (CD1). b2M is expressed on nearly all nucleated cells, with neurons being a notable exception. Circulating b2M is generated during normal HLA turnover. It can also dissociate from the MHC complex and circulate as full length and N-terminal truncated peptides of 93, 91, and 90 amino acids. It has been measured in a variety of body fluids, including serum, plasma, saliva, CSF, and urine. b2M freely passes through the glomerular membrane, but it is 99% actively reabsorbed and degraded in the proximal tubule cells. Circulating b2M levels are elevated in rheumatoid arthritis, systemic lupus erythematosus, viral infections, and conditions with decreased glomerular filtration. Human b2M shares 70% and 75% amino acid sequence identity with mouse and rat b2M, respectively.