

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Mcl-1 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 602901
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Mcl-1 Met231-Arg350 Accession # Q07820
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	8-25 µg/mL	See Below

DATA

Immunohistochemistry



Mcl-1 in Human Lymphoma. Mcl-1 was detected in immersion fixed paraffin-embedded sections of human lymphoma using Mouse Anti-Human Mcl-1 Monoclonal Antibody (Catalog # MAB8281) at 15 µg/mL overnight at 4 °C. Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counter-stained with hematoxylin (blue). Specific staining was localized to the cytoplasm and plasma membranes of lymphocytes. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.5 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Mcl-1 (myeloid cell leukemia-1) is a member of the Bcl-2 family of proteins. Alternative splicing creates two distinct isoforms. The 40 kDa, 350 amino acid (aa) isoform 1, also known as Mcl-1L (long), enhances cell survival by inhibiting apoptosis, while the ~35 kDa, 271 aa isoform 2, also known as Mcl-1S (short; with divergence in the last 41 aa) promotes apoptosis. The elimination of Mcl-1L is a required step for DNA damage-induced apoptosis. Mcl-1 can be modified by phosphorylation on S121 and T163 by JNK, which triggers apoptosis, or polyubiquitination, which enhances degradation. Within aa 231-350 (unique to Mcl-1L), human Mcl-1 shares 93% and 96% aa identity with mouse and rat Mcl-1, respectively.