

**DESCRIPTION**





<b>Species Reactivity</b>	Human/Mouse/Rat
<b>Specificity</b>	Detects human HHEX in direct ELISAs and Western blots.
<b>Source</b>	Recombinant Monoclonal Rabbit IgG Clone # 2018B
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human HHEX Thr111-Gly270 Accession # Q6UX06
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	See Below
<b>Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	See Below
<b>Immunocytochemistry</b>	2-25 µg/mL	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Hematopoietically-expressed homeobox protein (HHEX), also known as HEX, PRH and PRHX, is a 35-40 kDa member of the Homeobox family of transcription factors. Family members are distinguished by an evolutionarily conserved DNA-binding homeodomain of 60 amino acids (aa), which for HHEX spans aa 137-196. Human HHEX was initially isolated from hematopoietic tissue, and is present in several hematopoietic progenitors, where its expression is down-regulated during terminal cell differentiation. HHEX is also expressed in the anterior visceral endoderm during early mouse development, and in some adult tissues of endodermal origin, including liver, lung and thyroid. HHEX knockout in mice is embryonic lethal, with impaired forebrain, liver and thyroid development. Human HHEX is 270 aa in length, and over aa 111-270, shares 93% and 95% identity with mouse and rat HHEX, respectively.