

DESCRIPTION


| | |
|---------------------------|---|
| Species Reactivity | Human |
| Specificity | Detects human NFIL3/E4BP4 in direct ELISAs. |
| Source | Monoclonal Mouse IgG ₁ Clone # 714401 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | <i>E. coli</i> -derived recombinant human NFIL3/E4BP4 Lys140-Gly462 Accession # Q16649 |
| Formulation | Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|--|---------------|
| Flow Cytometry | 0.25 µg/mL | See Below |
| CyTOF-ready | Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation. | |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Reconstitution | Reconstitute at 0.5 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

NFIL3 (Nuclear Factor, Interleukin 3 Regulated), also called E4BP4 (adenovirus E4 promoter binding protein 4), is an approximately 58 kDa transcription factor of the bZIP (basic leucine zipper) family. It is a transcriptional regulator expressed in T and pro-B lymphocytes and cardiomyocytes, binds IL-3 and promoters as a homodimer, and enhances cell survival. NFIL3 has also been shown to be involved in the development of Innate Lymphoid Cells (ILCs; 1,2). The 462 amino acid (aa) human NFIL3 contains a basic DNA binding domain (aa 73-95), a leucine zipper (aa 96-124), and a transcriptional repression domain (aa 299-363) that is potentially regulated by phosphorylation at Ser301 and Ser353. Within the region used as an immunogen, human NFIL3 shares 83% aa sequence identity with mouse and rat NFIL3.

References:

1. Seillet C, et al (2014) J Exp Med. **211**:1733.
2. Geiger TL, et al (2014) J Exp Med. **211**:1723.