

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human NFIL3/E4BP4 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 714401
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human NFIL3/E4BP4 Lys140-Gly462 Accession # Q16649
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25 µg/mL	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

NFIL3 (Nuclear Factor, Interleukin 3 Regulated), also called E4BP4 (adenovirus E4 promoter binding protein 4), is an approximately 58 kDa transcription factor of the bZIP (basic leucine zipper) family. It is a transcriptional regulator expressed in T and pro-B lymphocytes and cardiomyocytes, binds IL-3 and promoters as a homodimer, and enhances cell survival. NFIL3 has also been shown to be involved in the development of Innate Lymphoid Cells (ILCs; 1,2). The 462 amino acid (aa) human NFIL3 contains a basic DNA binding domain (aa 73-95), a leucine zipper (aa 96-124), and a transcriptional repression domain (aa 299-363) that is potentially regulated by phosphorylation at Ser301 and Ser353. Within the region used as an immunogen, human NFIL3 shares 83% aa sequence identity with mouse and rat NFIL3.

References:

1. Seillet C, et al (2014) J Exp Med. **211**:1733.
2. Geiger TL, et al (2014) J Exp Med. **211**:1723.